arXiv:1412.3985v2 [math.OA] 29 Mar 2016

QUOTIENTS OF BANACH ALGEBRAS ACTING ON L^p-SPACES

EUSEBIO GARDELLA AND HANNES THIEL

Abstract. We show that the class of Banach algebras that can be isometrically represented on an L^p-space, for $p \neq 2$, is not closed under quotients. This answers a question asked by Le Merdy 20 years ago. Our methods are heavily reliant on our earlier study of Banach algebras generated by invertible isometries of L^p-spaces.

1. Introduction

An operator algebra is a closed subalgebra of the algebra $\mathcal{B}(\mathcal{H})$ of bounded linear operators on a Hilbert space $\mathcal{H}$. If A is an operator algebra and $I \subseteq A$ is a closed, two-sided ideal, then the quotient Banach algebra A/I is again an operator algebra, that is, it can be isometrically represented on a Hilbert space. This classical result is due to Lumer and Bernard, although the commutative case (when A is a uniform algebra) was proved earlier by Cole.

In Problem 3.8 of [LM96], Christian Le Merdy raised the question of whether this result can be generalized to Banach algebras acting on L^p-spaces for $p \in [1, \infty)$. More precisely, if $\mathscr{E}$ is a class of Banach spaces, we say that a Banach algebra A is an *$\mathscr{E}$-operator algebra* if there exist a Banach space E in $\mathscr{E}$ and an isometric homomorphism $\varphi\colon A \to \mathcal{B}(E)$. Given $p \in [1, \infty)$, we consider the class L^p of L^p-spaces, the class SL^p of Banach spaces that are isometrically isomorphic to subspaces of L^p-spaces, and the class QSL^p of Banach spaces that are quotients of SL^p-spaces.

The Bernard-Cole-Lumer Theorem asserts that L^2-operator algebras are closed under quotients. In Corollary 3.2 of [LM96], Le Merdy showed that QSL^p-operator algebras are closed under quotients. In Corollary 1.5.2.3 of [Jun96], Marius Junge showed the analogous result for SL^p-operator algebras. Since the classes QSL^2 and SL^2 both agree with L^2, the results of Le Merdy and Junge are generalizations of the Bernard-Cole-Lumer Theorem.

As the authors point out in [LM96] and [Jun96], the arguments used there are not suitable to deal with the more natural class of L^p-operator algebras. Indeed, the question of whether L^p-operator algebras are closed under quotients, for $p \neq 2$, remained open for 20 years. The case $p = 1$ of this question has recently been answered negatively in Theorem 6.2 of [GT14a], using a classical result of Malliavin on the failure of spectral synthesis for $\ell^1(\mathbb{Z})$. In this paper, we answer negatively the remaining cases of the question. (Even for $p = 1$, we construct new examples of quotients of $\ell^1(\mathbb{Z})$ which cannot be represented on an L^1-space. These are, in particular, semisimple, unlike those constructed in [GT14a].)

Date: 31 March 2016.

2010 *Mathematics Subject Classification.* Primary: 47L10, 43A15, Secondary: 46J10 .

Key words and phrases. Quotients of Banach algebras, L^p-space, Banach algebra of p-pseudofunctions.

The first named author was partially supported by the D. K. Harrison Prize from the University of Oregon. The second named author was partially supported by the Deutsche Forschungsgemeinschaft (SFB 878).

For $p \in [1,\infty)$, we consider the algebra $F^p(\mathbb{Z})$ of p-pseudofunctions on $\mathbb{Z}$. This algebra was introduced in the early 70's by Herz in [Her73], who denoted it $PF_p(\mathbb{Z})$. Algebras of p-pseudofunctions (also for locally compact groups other than $\mathbb{Z}$) have been studied in a number of places: [NR09]; [Phi13]; [GT14a]; [GT14c]; [GT14b], just to list a few.

The algebra $F^p(\mathbb{Z})$ is a semisimple, commutative Banach algebra with spectrum S^1. Given an open set V in S^1, we let I_V denote the largest closed two-sided ideal of $F^p(\mathbb{Z})$ that is supported on V. For $p \in [1,\infty) \setminus \{2\}$, and V neither empty nor dense in S^1, we show that $F^p(\mathbb{Z})/I_V$ is not an L^p-operator algebra; see Theorem 2.5. In fact, we even show that there is no injective, contractive homomorphism with closed range from $F^p(\mathbb{Z})/I_V$ to the algebra of bounded linear operators on *any* L^q-space for $q \in [1,\infty)$; see Remark 2.6.

Given the recent attention received by L^p-operator algebras, deciding whether these are closed under quotients becomes more relevant and technically useful. For example, consider the L^p-analogs $\mathcal{O}_n^p$ of the Cuntz algebras; see [Phi12]. These are all simple, and any contractive, non-zero representation of any of them on an L^p-space is automatically injective (in fact, isometric). For $p = 2$, these two properties are well-known to be equivalent. However, for $p \neq 2$, they are not, since quotients of L^p-operator algebras are not in general representable on L^p-spaces. These two properties of $\mathcal{O}_n^p$ therefore require separate and independent proofs. A similar problem arises with the L^p-analogs of irrational rotation algebras; see [GT15].

2. The example

A *representation* of a Banach algebra A on a Banach space E is a contractive homomorphism $\varphi\colon A \to \mathcal{B}(E)$. The representation is called *unital* (*injective*, *isometric*) if φ is.

We begin with some preparatory results. Our first lemma allows us to assume that a unital L^p-operator algebra has a unital, isometric representations on some L^p-space.

Lemma 2.1. *Let A be a unital Banach algebra and let $p \in [1,\infty)$. If A can be (injectively, isometrically) represented on an L^p-space, then there is a* unital *(injective, isometric) representation of A on an L^p-space.*

Proof. Let F be an L^p-space and let $\psi\colon A \to \mathcal{B}(F)$ be a contractive homomorphism. Then $e = \psi(1)$ is an idempotent with $\|e\| = 1$ in $\mathcal{B}(F)$. By Theorem 6 in [Tza69], the range E of e is an L^p-space. The cut-down homomorphism

$$\varphi\colon A \to \mathcal{B}(E) \cong \psi(1)\mathcal{B}(F)\psi(1)$$

is the desired map. It is clear that φ is injective or isomeric whenever ψ is, respectively. □

If A is a commutative unital Banach algebra, we will denote by $\Gamma_A\colon A \to C(\mathrm{Max}(A))$ its Gelfand transform, which is natural in the following sense. If $\varphi\colon A \to B$ is a unital homomorphism between commutative unital Banach algebras A and B, then the assignment $\mathrm{Max}(B) \to \mathrm{Max}(A)$ given by $I \mapsto \varphi^{-1}(I)$ defines a contractive homomorphism $\Gamma(\varphi)\colon C(\mathrm{Max}(A)) \to C(\mathrm{Max}(B))$ making the following diagram commute:

$$\begin{array}{ccc} A & \xrightarrow{\Gamma_A} & C(\mathrm{Max}(A)) \\ {\scriptstyle\varphi}\downarrow & & \downarrow{\scriptstyle\Gamma(\varphi)} \\ B & \xrightarrow[\Gamma_B]{} & C(\mathrm{Max}(B)). \end{array}$$

Remark 2.2. We recall the following fact about spectra of elements in Banach algebras. If B is a unital Banach algebra, A is a subalgebra containing the unit of B, and a is an element of A such that $\mathrm{sp}_A(a) = \partial\mathrm{sp}_A(a)$ as susbets of $\mathbb{C}$, then $\mathrm{sp}_A(a) = \mathrm{sp}_B(a)$. This can be deduced, for example, from Theorem in [Pal94]; see also the inclusions in the middle of page 219 there.

In particular, if the spectrum of an element of a Banach algebra is a subset of S^1, then it is equal to the spectrum computed in any unital algebra containing the given algebra.

Let $p \in [1, \infty)$. We first recall some facts about $F^p(\mathbb{Z})$; the reader is referred to [Phi13], [GT14c], and [GT14b] for more general versions of these statements, as well as their proofs.

The Banach algebra $F^p(\mathbb{Z})$ is a subalgebra of $\mathcal{B}(\ell^p(\mathbb{Z}))$ that can be defined as follows. Denote by $\lambda_p \colon \ell^1(\mathbb{Z}) \to \mathcal{B}(\ell^p(\mathbb{Z}))$ the left regular representation of $\mathbb{Z}$, which is given by $\lambda_p(f)\xi = f * \xi$ for $f \in \ell^1(\mathbb{Z})$ and $\xi \in \ell^p(\mathbb{Z})$. Then

$$F^p(\mathbb{Z}) = \overline{\lambda_p(\ell^1(\mathbb{Z}))} \subseteq \mathcal{B}(\ell^p(\mathbb{Z})).$$

The algebra $F^p(\mathbb{Z})$ also admits a description as a universal algebra with respect to invertible isometries on L^p-spaces. The algebra $F^p(\mathbb{Z})$ is clearly commutative, and its maximal ideal space is canonically homeomorphic to S^1. Moreover, its Gelfand transform is injective. For $p = 1$, there is a natural isometric identification $F^1(\mathbb{Z}) = \ell^1(\mathbb{Z})$, while for p and q in $[1, \infty)$, there is an abstract isometric isomorphism $F^p(\mathbb{Z}) \cong F^q(\mathbb{Z})$ if and only if p and q are either equal or (Hölder) conjugate; see Theorem 3.4 in [GT14b].

Notation 2.3. Let $p \in [1, \infty)$. We let Γ denote the Gelfand transform from $F^p(\mathbb{Z})$ to $C(S^1)$. For an open subset V of S^1, we set

$$I_V = \Gamma^{-1}(C_0(V)),$$

which is a closed, two-sided ideal in $F^p(\mathbb{Z})$. We will abbreviate $F^p(\mathbb{Z})$ to A, and the quotient $F^p(\mathbb{Z})/I_V$ to A_V. The Gelfand transform $\Gamma_{A_V} \colon A_V \to C(\mathrm{Max}(A_V))$ will be abbreviated to Γ_V.

Proposition 2.4. *Let $p \in [1, \infty)$, and let V be a nonempty open subset of S^1. We let $A = F^p(\mathbb{Z})$, I_V and A_V be as in Notation 2.3. Suppose that there exists $q \in [1, \infty)$ such that A_V is an L^q-operator algebra. Then the Gelfand transform $\Gamma_V \colon A_V \to C(S^1 \setminus V)$ is an isomorphism (although not necessarily isometric). In particular, and identifying $F^p(\mathbb{Z})$ with a subalgebra of $C(S^1)$ via Γ, it follows that every continuous function on $S^1 \setminus V$ is the restriction of a function in $F^p(\mathbb{Z})$.*

Proof. It is clear that $\mathrm{Max}(A_V)$ is canonically homeomorphic to $S^1 \setminus V$, so the range of Γ_V can be canonically identified with a subalgebra of $C(S^1 \setminus V)$. We claim that A_V is semisimple. Denote by $\pi \colon A \to A_V$ the canonical quotient map, and denote by $r \colon C(S^1) \to C(S^1 \setminus V)$ the restriction map. Observe that naturality of the Gelfand transform implies that the following diagram is commutative:

$$\begin{array}{ccc} A & \xrightarrow{\Gamma} & C(S^1) \\ {\scriptstyle\pi}\downarrow & & \downarrow{\scriptstyle r} \\ A_V & \xrightarrow[\Gamma_V]{} & C(S^1 \setminus V). \end{array} \tag{1}$$

It is a standard fact that semisimplicity of A_V is equivalent to injectivity of Γ_V. Let $a \in A_V$ satisfy $\Gamma_V(a) = 0$. Find $b \in A$ such that $\pi(b) = a$. Then $r(\Gamma(b)) = 0$, so $\Gamma(b)$ belongs to $\ker(r) = C_0(V)$. In other words, $b \in \Gamma^{-1}(C_0(V)) = I_V$. Hence $a = 0$ and A_V is semisimple.

It follows that there are natural identifications

$$A_V \cong \frac{\Gamma(F^p(\mathbb{Z}))}{\Gamma(F^p(\mathbb{Z})) \cap C_0(V)} \cong \frac{\Gamma(F^p(\mathbb{Z})) + C_0(V)}{C_0(V)}.$$

Observe that A_V is generated by the image $\pi(u)$ of the canonical generator u of $A = F^p(\mathbb{Z})$, which is an invertible isometry. Suppose that there exist $q \in [1, \infty)$, an L^q-space E, and an isometric representation $\varphi\colon A_V \to \mathcal{B}(E)$. By Lemma 2.1, we can assume that φ is unital. It is clear that $\varphi(\pi(u))$ generates $\varphi(A_V)$. Since φ is unital, $\varphi(\pi(u))$ is an invertible isometry of an L^q-space. Moreover, using Remark 2.2 at the first step, we have

$$\mathrm{sp}_{\mathcal{B}(E)}(\varphi(\pi(u))) = \mathrm{sp}_{\varphi(A_V)}(\varphi(\pi(u))) = \mathrm{sp}_{A_V}(\pi(u)) = S^1 \setminus V.$$

We claim that the Gelfand transform $\Gamma_{\varphi(A_V)}\colon \varphi(A_V) \to C(S^1 \setminus V)$ is an isomorphism. Once we show this, it will follow that Γ_V is also an isomorphism.

First, $\Gamma_{\varphi(A_V)}$ is clearly injective by semisimplicity of A_V. Suppose that $q = 2$. Then $\varphi(A_V)$ is a C^*-algebra, because it is generated by an invertible isometry of a Hilbert space (a unitary), and A_V is therefore self-adjoint. The claim is then an immediate consequence of Gelfand's theorem (and in this case $\Gamma_{\varphi(A_V)}$ is isometric). Assume now that $q \in [1, \infty) \setminus \{2\}$. In this case, and since the spectrum of $\varphi(\pi(u))$ in $\mathcal{B}(E)$ is not the whole circle, the result follows from part (1) of Corollary 5.21 in [GT14a]. The claim is proved, and the first part of the proposition follows.

For the second assertion in the statement of the proposition, consider the commutative diagram in (1). It follows that for every $f \in C(S^1 \setminus V)$, there exists $g \in A = F^p(\mathbb{Z})$ such that $\Gamma_V(\pi(g)) = f$. Regarding g as a function on S^1, this is equivalent to $g|_{S^1 \setminus V} = f$. □

Let $\theta \in \mathbb{R}$. Then it is easy to show that the homeomorphism $h_\theta\colon S^1 \to S^1$ given by $h_\theta(\zeta) = e^{2\pi i\theta}\zeta$ for $\zeta \in S^1$ induces an isometric automorphism of $F^p(\mathbb{Z})$. (We warn the reader that it is not in general true that any homeomorphism of S^1 induces an isometric, or even contractive, automorphism of $F^p(\mathbb{Z})$. In fact, when $p \neq 2$, the only homeomorphisms of S^1 that do so are the rotations and compositions of rotations with the homeomorphism $\zeta \mapsto \overline{\zeta}$ of S^1.)

The following is the main result of this paper. It asserts that for every $p \neq 2$, there is a quotient algebra of $F^p(\mathbb{Z})$ that has no isometric representation on any L^q-space for $q \in [1, \infty)$.

Theorem 2.5. *Let $p \in [1, \infty) \setminus \{2\}$, let V be a nonempty open subset of S^1. We let $A = F^p(\mathbb{Z})$, I_V and A_V be as in Notation 2.3. Assume that V is not dense in S^1. Then A_V cannot be isometrically represented on* any *L^q-space for $q \in [1, \infty)$.*

Proof. We argue by contradiction, so let V be an open subset of S^1 as in the statement, and suppose that there exists $q \in [1, \infty)$ such that A_V can be isometrically represented on an L^q-space.

Let $f \in C(S^1)$. We claim that f belongs to $\Gamma(F^p(\mathbb{Z}))$. Once we prove this, it will follow from part (2) of Corollary 3.20 in [GT14c] that $p = 2$, and hence the proof will be complete.

Let W be a nonempty open subset of S^1 such that $V \cap W = \emptyset$. With the notation used in the comments before this theorem, and using compactness of S^1, find $n \in \mathbb{N}$ and $\theta_1, \ldots, \theta_n \in \mathbb{R}$ such that $\bigcup_{j=1}^{n} h_{\theta_j}(W) = S^1$. For $j \in \{1, \ldots, n\}$, set $V_j = h_{\theta_j}(V)$ and $W_j = h_{\theta_j}(W)$. There is an isometric isomorphism

$$A_{h_{\theta_j}(V)} \cong A_V,$$

so the Banach algebra $A_{h_{\theta_j}(V)}$ can be isomorphically represented on an L^q-space. It follows from Proposition 2.4 that every continuous function on $S^1 \setminus V_j$ is in the image of Γ_{V_j}. In particular, every continuous function on $\overline{W_j}$ is in the image of Γ_{V_j}.

From now on, we identify the algebras $A, A_{V_1}, \ldots, A_{V_n}$ with their images under their Gelfand transforms. In particular, for $j = 1, \ldots, n$, every continuous function on $\overline{W_j}$ is the restriction of a function in A.

Choose continuous functions $k_1, \ldots, k_n \colon S^1 \to \mathbb{R}$ satisfying

(1) $0 \leq k_j \leq 1$ for $j = 1, \ldots, n$;
(2) $\operatorname{supp}(k_j) \subseteq W_j$ for $j = 1, \ldots, n$;
(3) $\sum_{j=1}^{n} k_j(\zeta) = 1$ for all $\zeta \in S^1$;
(4) k_j belongs to $F^p(\mathbb{Z})$ for $j = 1, \ldots, n$ (for example, take $k_j \in C^\infty(S^1)$).

For $j = 1, \ldots, n$, choose a function $g_j \in F^p(\mathbb{Z})$ such that $(g_j)|_{W_j} = f|_{W_j}$. Then the product $g_j k_j$ belongs to $F^p(\mathbb{Z})$ because each of the factors does. Since the support of k_j is contained in W_j, and f and g_j agree on W_j, we have $f k_j = g_j k_j$ for $j = 1, \ldots, n$. Now,

$$f = f \cdot \left(\sum_{j=1}^{n} k_j \right) = \sum_{j=1}^{n} g_j k_j,$$

so f belongs to $F^p(\mathbb{Z})$, and the claim is proved.

We have shown that Gelfand transform $\Gamma \colon F^p(\mathbb{Z}) \to C(S^1)$ is surjective. Since $F^2(\mathbb{Z})$ is canonically isomorphic to $C(S^1)$, we must have $p = 2$ by part (2) of Corollary 3.20 in [GT14c]. This is a contradiction, and the result follows. □

Remark 2.6. The same argument as in Proposition 2.4 and Theorem 2.5 shows that if $V \subseteq S^1$ is a non-trivial, non-dense open subset, then there is no contractive, injective representation of A_V on any L^q-space *with closed range*. On the other hand, there certainly are contractive, injective representations of A_V on L^q-spaces for every $q \in [1, \infty)$. Indeed, for $q \in [1, \infty)$, any commutative semisimple Banach algebra B can be represented on an L^q-space (although rarely can it be represented with closed range). Indeed, $C_0(\operatorname{Max}(B))$ is isometrically represented on an L^q-space for every q, and such a representation can be composed with the Gelfand transform to get the desired contractive representation of B on an L^q-space.

Remark 2.7. In contrast to Theorem 2.5, some (non-trivial) quotients of $F^p(\mathbb{Z})$ are isometrically representable on L^p-spaces. For example, if V is the complement of the set of n-th roots of unity in S^1 for some $n \in \mathbb{N}$, then $F^p(\mathbb{Z})/I_V$ is canonically isometrically isomorphic to $F^p(\mathbb{Z}_n)$. (This identification is induced by the quotient map $\mathbb{Z} \to \mathbb{Z}_n$.) An analogous statement holds for the translates of V. We do not know, however, whether these are the only quotients of $F^p(\mathbb{Z})$ that can be represented on L^p-spaces. We therefore suggest:

Problem 2.8. Characterize those ideals I of $F^p(\mathbb{Z})$ such that $F^p(\mathbb{Z})/I$ can be isometrically represented on an L^p-space.

We do not know whether $F^p(\mathbb{Z})$ has spectral synthesis, except for $p = 1$ (in which case it does not) and $p = 2$ (in which case it does). Since Banach algebras generated by an invertible isometry of an L^p-space together with its inverse are automatically semisimple by the results in [GT14a], we conclude that for $F^p(\mathbb{Z})/I$ to be isometrically representable on an L^p-space, there must exist an open subset $V \subseteq S^1$ such that $I = I_V$ (and V must be dense by Theorem 2.5). This means that Problem 2.8 can be solved without knowing whether $F^p(\mathbb{Z})$ has spectral synthesis, that is, without knowing whether every ideal of $F^p(\mathbb{Z})$ is of the form I_V.

We do not know whether density of V is sufficient for $F^p(\mathbb{Z})/I_V$ to be representable on an L^p-space.

We conclude this paper with an observation. If A is a Banach algebra and $a \in A$, we denote by $B(a)$ the smallest Banach subalgebra of A containing a.

Remark 2.9. Suppose that p is not an even integer, and let $V \subseteq S^1$ be a nonempty, non-dense open subset. We let $A = F^p(\mathbb{Z})$, I_V and A_V be as in Notation 2.3. It follows from Corollary 1.5.2.3 in [Jun96] that A_V is an SL^p-operator algebra, so there exists an invertible isometry v of an SL^p-space E such that $B(u)$ is isometrically isomorphic to A_V. By Theorem I in [Rud76], there exist a L^p-space F containing E as a closed subspace, and a canonical invertible isometry w of F extending v. By naturality of the construction, one may be tempted to guess that $B(v)$ and $B(w)$ are isometrically isomorphic. However, Theorem 2.5 shows that this is not the case.

References

[GT14a] E. Gardella and H. Thiel, Banach algebras generated by an invertible isometry of an L^p-space, J. Funct. Anal. **269** (2015), 1796–1839.

[GT14b] E. Gardella and H. Thiel, Functoriality of group algebras acting on L^p-spaces, preprint (arXiv:1408.6137 [math.FA]), 2014.

[GT14c] E. Gardella and H. Thiel, Group algebras acting on L^p-spaces, J. Fourier Anal. Appl. **21** (2015), no. 6, 1310–1343.

[GT15] E. Gardella and H. Thiel, Analogs of irrational rotation algebras on L^p-spaces, in preparation, 2015.

[Her73] C. Herz, Harmonic synthesis for subgroups, *Ann. Inst. Fourier (Grenoble)* **23** (1973), 91–123. MR 0355482 (50 #7956). Zbl 0257.43007.

[Jun96] M. Junge, Factorization theory for spaces of operators, Habilitationsschrift, 1996.

[LM96] C. Le Merdy, Representation of a quotient of a subalgebra of $B(X)$, *Math. Proc. Cambridge Philos. Soc.* **119** (1996), 83–90. MR 1356160 (97e:47074). Zbl 0847.46029.

[NR09] M. Neufang and V. Runde, Column and row operator spaces over QSL_p-spaces and their use in abstract harmonic analysis, *J. Math. Anal. Appl.* **349** (2009), 21–29. MR 2455727 (2009i:46105). Zbl 1155.43003.

[Pal94] Theodore W. Palmer, Banach algebras and the general theory of C^*-algebras. Vol. I. Algebras and Banach algebras. Encyclopedia of Mathematics and its Applications, **49**. Cambridge University Press, Cambridge, 1994. xii+794 pp.

[Phi12] N. C. Phillips, Analogs of Cuntz algebras on L^p spaces, preprint (arXiv:1201.4196 [math.FA]), 2012.

[Phi13] N. C. Phillips, Crossed products of L^p operator algebras and the K-theory of Cuntz algebras on L^p spaces, preprint (arXiv:1309.6406 [math.FA]), 2013.

[Rud76] W. Rudin, L^p-isometries and equimeasurability, *Indiana Univ. Math. J.* **25** (1976), 215–228. MR 0410355 (53 #14105). Zbl 0326.46011.

[Tza69] L. Tzafriri, Remarks on contractive projections in L_p-spaces, *Israel J. Math.* **7** (1969), 9–15. MR 0248514 (40 #1766). Zbl 0184.15103.

Eusebio Gardella Mathematisches Institut, Fachbereich Mathematik und Informatik der Universität Münster, Einsteinstrasse 62, 48149 Münster, Germany.

E-mail address: gardella@uni-muenster.de

Hannes Thiel Mathematisches Institut, Fachbereich Mathematik und Informatik der Universität Münster, Einsteinstrasse 62, 48149 Münster, Germany.

E-mail address: hannes.thiel@uni-muenster.de

URL: http://wwwmath.uni-muenster.de/u/hannes.thiel/